

MassTag Polymerase Chain Reaction for Differential Diagnosis of Viral Hemorrhagic Fevers

**Gustavo Palacios,*¹ Thomas Briese,*¹
Vishal Kapoor,* Omar Jabado,* Zhiqiang Liu,*
Marietjie Venter,† Junhui Zhai,* Neil Renwick,*
Allen Grolla,‡ Thomas W. Geisbert,§
Christian Drosten,¶ Jonathan Towner,#
Jingyue Ju,* Janusz Paweska,**
Stuart T. Nichol,# Robert Swanepoel,**
Heinz Feldmann,††† Peter B. Jahrling,‡‡
and W. Ian Lipkin***

Viral hemorrhagic fevers are associated with high rates of illness and death. Although therapeutic options are limited, early differential diagnosis has implications for containment and may aid in clinical management. We describe a diagnostic system for rapid, multiplex polymerase chain reaction identification of 10 different causes of viral hemorrhagic fevers.

Increasing international travel, trafficking in wildlife, political instability, and terrorism have made emerging infectious diseases a global concern. Viral hemorrhagic fevers (VHF) warrant specific emphasis because of their high rates of illness and death, and the potential for rapid dissemination by human-to-human transmission. The term “viral hemorrhagic fever” characterizes a severe multisystem syndrome associated with fever, shock, and bleeding diathesis caused by infection with any of several RNA viruses, including Ebola virus and Marburg virus (MARV) (family *Filoviridae*); Lassa virus (LASV) and the South American hemorrhagic fever viruses Guanarito virus, Junín

virus, Machupo virus, and Sabiá virus (*Arenaviridae*); Rift Valley fever virus (RVFV), Crimean-Congo hemorrhagic fever virus (CCHFV), and hantaviruses (*Bunyaviridae*); and Kyasanur Forest disease virus (KFDV), Omsk hemorrhagic fever virus, yellow fever virus (YFV), and dengue viruses (*Flaviviridae*) (1,2). Although clinical management of VHF is primarily supportive, early diagnosis is needed to contain the contagion and implement public health measures, especially if agents are encountered out of their natural geographic context.

Vaccines have been developed for YFV, RVFV, Junín virus, KFDV, and hantaviruses (3–7), but only YFV vaccine is widely available. Early treatment with immune plasma was effective in Junín virus infection (8). The nucleoside analog ribavirin may be helpful if given early in the course of Lassa fever (9), Crimean-Congo hemorrhagic fever (10), or hemorrhagic fever with renal syndrome (11) and is recommended in postexposure prophylaxis and early treatment of arenavirus and bunyavirus infections (12).

Methods for direct detection of nucleic acids of microbial pathogens in clinical specimens are rapid, sensitive, and obviate the need for high-level biocontainment. Numerous systems are described for nucleic acid detection of VHF agents; however, none are multiplex (13). Although geographic location or travel history of suspected patients usually restricts the number of agents to be considered, diagnosis of VHF may be difficult in case of an intentional release (12). Symptoms of VHF are initially nonspecific and may include fever, headache, myalgia, and gastrointestinal or upper respiratory tract complaints (1); thus, assays that allow simultaneous consideration of multiple agents are needed.

We recently described the application of MassTag polymerase chain reaction (PCR) in the context of differential diagnosis of respiratory disease (14). MassTag PCR is a multiplex assay in which microbial gene targets are coded by a library of 64 distinct mass tags. Nucleic acids (RNA or DNA) are amplified by multiplex (reverse transcription–) PCR using up to 64 primers, each labeled by a photo-cleavable link with a different molecular weight tag. After separation of the amplification products from unincorporated primers and release of the mass tags from the amplicons by UV irradiation, tag identity is analyzed by mass spectrometry. The identity of the microbe in the clinical sample is determined by the presence of its 2 cognate tags, 1 from each primer.

The Study

To facilitate rapid differential diagnosis of VHF agents, we established the Greene MassTag Panel VHF version 1.0, which comprises the following targets: Ebola Zaire virus

*Columbia University, New York, New York, USA; †University of Pretoria and National Health Laboratory Services, Pretoria, South Africa; ‡Public Health Agency of Canada, Winnipeg, Manitoba, Canada; §United States Army Medical Research Institute of Infectious Diseases, Fort Detrick, Frederick, Maryland, USA; ¶Bernhard-Nocht-Institute of Tropical Medicine, Hamburg, Germany; #Centers for Disease Control and Prevention, Atlanta, Georgia, USA; **National Institute for Communicable Diseases, Sandringham, South Africa; ††University of Manitoba, Winnipeg, Manitoba, Canada; and ‡‡National Institutes of Allergy and Infectious Diseases Integrated Research Facility, Fort Detrick, Frederick, Maryland, USA

¹These authors contributed equally to this article.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 1 APR 2006		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE MassTag polymerase chain reaction for differential diagnosis of viral hemorrhagic fever, Emerging Infectious Diseases 12:692 - 695				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Palacios, G Briese, T Kapoor, V Jabado, O Liu, Z Venter, M Zhai, J Renwick, N Grolla, A Geisbert, TW Drosten, C Towner, J Ju, J Paweska, J Nichol, ST Swanepoel, R Feldmann, H Jahrling, PB Lipkin, WI				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) United States Army Medical Research Institute of Infectious Diseases, Fort Detrick, MD				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Viral hemorrhagic fevers are associated with high rates of illness and death. Although therapeutic options are limited, early differential diagnosis has implications for containment and may aid in clinical management. We describe a diagnostic system for rapid, multiplex polymerase chain reaction identification of 10 different causes of viral hemorrhagic fevers.					
15. SUBJECT TERMS methods, PCR, MassTag, differential diagnosis, hemorrhagic fever, viral					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

(ZEBOV), Ebola Sudan virus (SEBOV), MARV, LASV, RVFV, CCHFV, Hantaan virus (HNTV), Seoul virus (SEOV), YFV, and KFDV. Oligonucleotide primers were designed in conserved genomic regions to detect the broadest number of members for a given pathogen species. We developed a software program that culls sequence information from GenBank, performs multiple alignments with ClustalW, and designs primers optimized for multiplex PCR. The program uses a greedy algorithm to identify conserved sequences and create the minimum set of primers for amplification of all sequences in the alignment. Primers are selected within standard design constraints whenever possible (melting temperature 55°C–65°C, guanine-cytosine content 40%–60%, no hairpins); degenerate positions are introduced in cases where template divergence requires more flexibility. Although degeneracy is not tolerated in the five 3' nucleotides, MassTag PCR allows up to 4 nonneighboring variable positions per primer. Primers are checked by the basic local alignment search tool for potential hybridization to sequenced vertebrate genomes (Table 1).

Because only released mass tags are analyzed, staggering the size of amplification products created in multiplex reactions is unnecessary; thus, primers are selected for efficient and consistent performance irrespective of amplicon size (typically 80–200 bp). Before committing to synthesis of tagged primers, the functionality of candidate multiplex primer panels is examined in a series of amplification reactions that use prototype templates representing individual microbial targets. Primers that fail to yield a single, specific product band in agarose gel analysis are

replaced. Target sequence standards for evaluation are cloned into pCR2.1-TOPO (Invitrogen, Carlsbad, CA, USA) by using PCR amplification of cDNA templates obtained by reverse transcription (RT) of extracts from infected, cultured cells or by assembly of overlapping synthetic polynucleotides.

The agents assayed in the VHF panel have RNA genomes; thus, assay sensitivity was determined by using synthetic RNA standards. Synthetic RNA standards were generated from linearized target sequence plasmids by using T7 polymerase (mMessage mMachine, Invitrogen). After quantitation by UV spectrometry, RNA was serially diluted in 2.5 µg/mL yeast tRNA (Sigma, St. Louis, MO, USA), reverse transcribed with random hexamers by using Superscript II (Invitrogen), and analyzed by MassTag PCR as previously described (14). QIAquick 96 PCR purification cartridges (Qiagen, Hilden, Germany, with modified binding and wash buffers) were used to remove unincorporated primers before tags were decoupled from amplification products by UV photolysis in a flow cell and analyzed in a quadrupole mass spectrometer by using positive-mode atmospheric pressure chemical ionization (APCI-MS, Agilent Technologies, Palo Alto, CA, USA). The sensitivity of the 10-plex VHF panel with synthetic RNA standards was ≤50 RNA copies per assay (Table 2). Sensitivity and specificity of multiplex primer panels is assessed empirically by using calibrated synthetic standards as well as tissue culture–derived viral nucleic acid for each assembled panel.

Tissue culture extracts were used to examine assay specificity. Random primed cDNA obtained from cultures

Table 1. Greene MassTag panel VHF version 1.0*

Target	MassTag	Name	Sequence	Gene
ZEBOV	718 (fwd)	EboZA-U234	AACACCGGGTCTTAATTCTTATATCAA	L
	646 (rev)	EboZA-L319	GGTGGTAAATTCCTAGTAGTTCTTT	
SEBOV	503 (fwd)	EboSU-U416	CGAGCCTAACGTTTTGGGC	L
	630 (rev)	EboSU-L489	GCTCCAGGAATTGTTCCGGTA	
MARV	654 (fwd)	MARV-U12816C	CCCTCCATATCTTAGACAACATATTGTG	L
	395 (rev)	MARV-L12994	CCCAACACTCCTGGTTCACAGC	
LASV†	558 (fwd)	Las4-U92	ACTGCATTYTCACTTYCTRGAATC	NP
	686 (rev)	Las4-L257	CCRGGYTTGACCACTGCTGT	
RVFV	658 (fwd)	RVF-U578	GGATTGACCTGTGCCTGTTGC	N
	495 (rev)	RVF-L660	GCATTAGAAATGTCCTCTTTTGCTGC	
CCHFV	499 (fwd)	CCHV-U4	AGAAACACGTGCCGCTTACGCCCA	N
	710 (rev)	CCHV-L120	CCATTTCCYTTYTTTAACTCYTCAAACCA	
HNTV	479 (fwd)	HAN-U179	AYACAGCAGCAGTTAGCCTCCT	N
	702 (rev)	HAN-L245	GCT GCC GTA RGT AGT CCC TGTT	
SEOV	455 (fwd)	SEO-U243	CAGGATTGCAGCAGGGAAGA	N
	602 (rev)	SEOUL-L309	ATGATCACCAGGYTCTACCCC	
YFV	467 (fwd)	YF-U186	GCTGGGAGCGCGGTATC	NS5
	670 (rev)	YF-L249	GGAAGCCCAATGGTCCTCAT	
KFDV	483 (fwd)	KYF-U170	TGGAAGCCTGGCTGAAAGAG	NS5
	614 (rev)	KYF-L233	TCATCCCCACTGACCAGCAT	

*ZEBOV, Ebola Zaire virus; SEBOV, Ebola Sudan virus; MARV, Marburg virus; LASV, Lassa virus; RVFV, Rift Valley fever virus; CCHFV, Crimean-Congo hemorrhagic fever virus; HNTV, Hantaan virus; SEOV, Seoul virus; YFV, yellow fever virus; KFDV, Kyasanur Forest disease virus; fwd, forward; rev, reverse.

†Primers were designed on Lassa lineage IV sequences (15) and the recently identified outlier sequence Lassa AV (AF256121).

DISPATCHES

Table 2. Sensitivity of detection with synthetic RNA standards

Pathogen*	Detection threshold (RNA copies)†
ZEBOV	20
SEBOV	20
MARV	20
LASV	20
RVFV	20
CCHFV	50
HNTV	20
SEOV	50
YFV	20
KFDV	20

*ZEBOV, Ebola Zaire virus; SEBOV, Ebola Sudan virus; MARV, Marburg virus; LASV, Lassa virus; RVFV, Rift Valley fever virus; CCHFV, Crimean-Congo hemorrhagic fever virus; HNTV, Hantaan virus; SEOV, Seoul virus; YFV, yellow fever virus; KFDV, Kyasanur Forest disease virus.

†RNA copies refers to the number of molecules subjected to reverse transcription; half of the reverse transcription reaction was then used for polymerase chain reaction amplification.

of ZEBOV, SEBOV, MARV, YFV isolates from the Gambia and Côte d'Ivoire, RVFV, CCHFV, HNTV, SEOV, and LASV strains Josiah, NL, and AV were subjected to mass tag analysis. In all instances, only the appropriate cognate mass tags were detected (data not shown). No spurious signal was identified in assays with water or RNA controls.

Performance with clinical materials was tested by using blood, sera, or oral swabs from 24 human patients of VHF

previously diagnosed through virus isolation, RT-PCR, or antigen detection enzyme-linked immunosorbent assay. Differential diagnosis by blinded MassTag PCR analysis was accurate in all cases (Table 3). For the samples from the 2005 Angola Marburg outbreak the result of MassTag PCR was similar to that of diagnostic single-plex PCR. ZEBOV sample 5004, obtained on day 17 of illness when serologic test results were positive for immunoglobulin M (IgM) and IgG, was negative by viral culture but positive in MassTag PCR.

Conclusions

These results confirm earlier work in respiratory diseases that show that MassTag PCR offers a rapid, sensitive, specific, and economic approach to differential diagnosis of infectious diseases. Small, low-cost, or mobile APCI-MS units extend the applicability of this technique beyond selected reference laboratories. Given the capacity of the method to code for up to 32 genetic targets, we are expanding the hemorrhagic fever panel to include additional viruses (dengue and South American hemorrhagic fever viruses) and are exploring the inclusion of bacterial and parasitic agents that may result in similar clinical signs and symptoms and, thus, have to be considered in differential diagnosis.

Table 3. MassTag polymerase chain reaction analysis of clinical specimens from viral hemorrhagic fever patients*

Previous diagnosis	Sample identification	Sample type	Year/origin	MassTag result†
ZEBOV	5015	Serum	1995/Kikwit, DRC	+++ , ZEBOV
ZEBOV	5014	Serum	1995/Kikwit, DRC	+++ , ZEBOV
ZEBOV	5004	Serum	1995/Kikwit, DRC	+++ , ZEBOV
ZEBOV	6317	Serum	1995/Kikwit, DRC	+++ , ZEBOV
ZEBOV	6313	Serum	1995/Kikwit, DRC	+++ , ZEBOV
MARV	246-00-5	Hemolyzed whole blood	2000/Durba, DRC	+, MARV
MARV	226-00-4	Hemolyzed whole blood	2000/Durba, DRC	++, MARV
MARV	246-00-7	Hemolyzed whole blood	2000/Durba, DRC	+, MARV
MARV	98-00-2	Hemolyzed whole blood	2000/Durba, DRC	+++ , MARV
MARV	461	Blood	2005/Uige, Angola	+++ , MARV
MARV	462	Oral swab	2005/Uige, Angola	+++ , MARV
MARV	475	Blood	2005/Uige, Angola	++, MARV
MARV	476	Oral swab	2005/Uige, Angola	+, MARV
LASV	98-04-1	Serum	2004/Sierra Leone	+++ , LASV
LASV	98-04	Serum	2004/Sierra Leone	++, LASV
LASV	98-04-5	Serum	2004/Sierra Leone	+, LASV
LASV	80-04-1	Serum	2004/Sierra Leone	+++ , LASV
RVFV	98002009	Serum	1998/Kenya	+, RVFV
RVFV	H6061989	Serum	1998/Kenya	+, RVFV
RVFV	98002019	Serum	1998/Kenya	++, RVFV
RVFV	77-04	Serum	2004/Namibia	++, RVFV
CCHFV	187-86	Serum	1986/South Africa	+, CCHFV
CCHFV	30-93	Serum	1993/South Africa	+++ , CCHFV
CCHFV	465-88	Serum	1988/South Africa	+++ , CCHFV
CCHFV	407-89	Serum	1989/South Africa	+++ , CCHFV
CCHFV	215-90	Serum	1990/South Africa	++, CCHFV

*ZEBOV, Ebola Zaire virus; MARV, Marburg virus; LASV, Lassa virus; RVFV, Rift Valley fever virus; CCHFV, Crimean-Congo hemorrhagic fever virus; DRC, Democratic Republic of Congo.

†Relative ranking of results: +, signal-to-noise ratio ≤ 4 ; ++, signal-to-noise ratio >4 and <8 ; +++, signal-to-noise ratio ≥ 8 .

This work was supported by National Institutes of Health awards AI51292, AI056118, AI55466, and U54AI57158 (Northeast Biodefense Center-Lipkin) and the Ellison Medical Foundation.

Dr Palacios is an associate research scientist in the Jerome L. and Dawn Greene Infectious Disease Laboratory at the Columbia University Mailman School of Public Health. His research focuses on the molecular epidemiology of viruses, virus interactions with their hosts, and innovative pathogen detection methods.

References

- Peters CJ, Zaki SR. Role of the endothelium in viral hemorrhagic fevers. *Crit Care Med.* 2002;30(5 Suppl):S268–73.
- Geisbert TW, Jahrling PB. Exotic emerging viral diseases: progress and challenges. *Nat Med.* 2004;10(12 Suppl):S110–21.
- Pugachev KV, Guirakhoo F, Monath TP. New developments in flavivirus vaccines with special attention to yellow fever. *Curr Opin Infect Dis.* 2005;18:387–94.
- Pittman PR, Liu CT, Cannon TL, Makuch RS, Mangiafico JA, Gibbs PH, et al. Immunogenicity of an inactivated Rift Valley fever vaccine in humans: a 12-year experience. *Vaccine.* 1999;18:181–9.
- Enria DA, Barrera Oro JG. Junin virus vaccines. *Curr Top Microbiol Immunol.* 2002;263:239–61.
- Hooper JW, Li D. Vaccines against hantaviruses. *Curr Top Microbiol Immunol.* 2001;256:171–91.
- Dandawate CN, Desai GB, Achar TR, Banerjee K. Field evaluation of formalin inactivated Kyasanur forest disease virus tissue culture vaccine in three districts of Karnataka state. *Indian J Med Res.* 1994;99:152–8.
- Enria DA, Maiztegui JI. Antiviral treatment of Argentine hemorrhagic fever. *Antiviral Res.* 1994;23:23–31.
- McCormick JB, King JJ, Webb PA, Scribner CL, Craven RB, Johnson KM, et al. Lassa fever. Effective therapy with ribavirin. *N Engl J Med.* 1986;314:20–6.
- Ozkurt Z, Kiki I, Erol S, Erdem F, Yilmaz N, Parlak M, et al. Crimean-Congo hemorrhagic fever in eastern Turkey: clinical features, risk factors and efficacy of ribavirin therapy. *J Infect.* Epub 2005 Jun 13.
- Huggins JW, Hsiang CM, Cosgriff TM, Guang MY, Smith JI, Wu ZO, et al. Prospective, double-blind, concurrent, placebo-controlled clinical trial of intravenous ribavirin therapy of hemorrhagic fever with renal syndrome. *J Infect Dis.* 1991;164:1119–27.
- Borio L, Inglesby T, Peters CJ, Schmaljohn AL, Hughes JM, Jahrling PB, et al. Hemorrhagic fever viruses as biological weapons: medical and public health management. *JAMA.* 2002;287:2391–405.
- Drosten C, Kummerer BM, Schmitz H, Gunther S. Molecular diagnostics of viral hemorrhagic fevers. *Antiviral Res.* 2003;57:61–87.
- Briese T, Palacios G, Kokoris M, Jabado O, Liu Z, Renwick N, et al. Diagnostic system for rapid and sensitive differential detection of pathogens. *Emerg Infect Dis.* 2005;11:310–3.
- Bowen MD, Rollin PE, Ksiazek TG, Hustad HL, Bausch DG, Demby AH, et al. Genetic diversity among Lassa virus strains. *J Virol.* 2000;74:6992–7004.

Address for correspondence: Thomas Briese, Jerome L. and Dawn Greene Infectious Disease Laboratory, Mailman School of Public Health, Columbia University, 722 W 168th St, Rm 1801, New York, NY 10032, USA; fax: 212-342-9044; email: thomas.briese@columbia.edu

Use of trade names is for identification only and does not imply endorsement by the Public Health Service or by the U.S. Department of Health and Human Services.

EMERGING INFECTIOUS DISEASES

Full text free online at
www.cdc.gov/eid

The print journal is available at no charge to public health professionals

☐ YES, I would like to receive Emerging Infectious Diseases.

Please print your name and business address in the box and return by fax to 404-639-1954 or mail to

EID Editor
CDC/NCID/MS D61
1600 Clifton Road, NE
Atlanta, GA 30333

Moving? Please give us your new address (in the box) and print the number of your old mailing label here _____

Copyright of Emerging Infectious Diseases is the property of Centers for Disease Control & Prevention (CDC) and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.